Opioid Overdose Response

Follow these steps for naloxone ampoule injection

1

SAFETY

Check the person and area for anything unsafe.



2

OPIOID OVERDOSE? LOOK FOR THE SIGNS

- Not responding
- Pinned pupils
- Not breathing normally shallow slow breathing or snoring
- Pale or cold skin, blue or grey lips and fingertips



3

CALL TRIPLE ZERO (000)

I think it's an overdose



4

LIE PERSON ON SIDE

Put person in recovery position.



5


ATTACH THE NEEDLE TO A SYRINGE



6

SWIRL THE AMPOULE


Tap or swirl ampoule so liquid is at the base.



7

HOLD THE AMPOULE


Hold base of ampoule in one hand and pinch the top with the other.




continued on next page











ARE THEY BREATHING?

If not, use *rescue*breathing if you know
how – lie person on back
and give 2 breaths.



RECOVERY


Put person in the recovery position.
Stay with them until Ambulance arrives.



14

REPEAT DOSE IF NEEDED

If no response, inject another dose every 2 to 3 minutes until person recovers or ambulance arrives.



Use naloxone to reverse opioid overdose (e.g. heroin, morphine, methadone, oxycodone, prescription opioids, fentanyl). It takes 2–5 minutes to start working and lasts about 30-90 minutes.

To receive this document in another format email <aod.enquiries@health.vic.gov.au>

Authorised and published by the Victorian Government, 1 Treasury Place, Melbourne. © State of Victoria, Australia, Department of Health. ISBN 978-1-76131-461-2 (pdf/online/MS word).